



RIVER VALLEY SCHOOL DISTRICT

660 West Daley Street

≈

Spring Green, Wisconsin 53588

≈

Phone: 608-588-2551

428/528 Exhibit

Periods of Exclusion from School with Infectious Conditions

Personal hygiene measures such as hand washing, covering the mouth and nose when coughing or sneezing, covering weeping sores, not sharing food or drinks, and not attending school when ill or suffering from diarrhea are important means of limiting the transmission of a number of common infectious conditions.

The student/staff who have been in contact with the following conditions, to be excluded from school for the periods specified.

For the purposes of these guidelines, communicable diseases include, but are not limited to, the following:

Condition	Criteria for Exclusion from School	Onsite control and prevention measures
Mumps	Exclude for 5 days after swelling onset.	Provide immunization records for exposed individuals to public health officials Good handwashing and hygiene. Avoid kissing, sharing drinks, or utensils, use proper disinfection of surfaces and toys.
Pink Eye	None, unless fever, behavior change, or unable to avoid touching the eye. Antibiotics not required for return	Good handwashing and hygiene. Avoid kissing, sharing drinks, or utensils, use proper disinfection of surfaces and toys.
Strep Throat	Exclude for 24 hours after initiation of appropriate antibiotic and fever resolved. Fever free without fever reducing medication for 24 hours	Avoid kissing, sharing drinks, or utensils; exclude infected food handlers; minimize contact with respiratory and oral secretions. Good handwashing and hygiene.
Influenza	Exclude until fever resolved for 24 hours Fever free without fever reducing	Handwashing and good personal hygiene including covering coughs and sneezes.

	medication for 24 hours	
Pertussis (whooping cough)	Exclude until after 5 days of antibiotic treatment; if no antibiotic treatment, exclude 21 days after cough onset	Refer symptomatic individuals to health care provider for evaluation. Handwashing and good personal hygiene including covering coughs and sneezes
Diarrhea	Exclude until asymptomatic for 24 hours.	3 or more loose stools in 24 hours. Outbreaks of diarrhea in group settings are reportable to the public health department. Good handwashing and hygiene; proper disinfection of tables, toys, and food preparation areas.
Gastroenteritis (viral)	Exclude until asymptomatic for 24 hours for rotavirus and 48 hours with norovirus	Good handwashing and hygiene; proper disinfection of tables, toys, and food preparation areas. Alcohol based hand products are not effective. Soap and water must be used.
Hepatitis A Virus	Exclude for 14 days after onset of symptoms or 10 days after onset of jaundice.	Consider Hepatitis A vaccine. Exposed staff should not prepare meals for others. Good handwashing and hygiene; proper disinfection of tables, toys, and food preparation areas.
Pinworm Infection	None, unless has diarrhea.	Frequent, good handwashing, particularly by infected child and staff assisting in toileting; keep fingernails clean and short; prevent fingers in mouth.
Clostridium Difficile Infection (c.diff)	Exclude until asymptomatic for 48 hours.	Good handwashing and hygiene; proper disinfection of tables, toys, and food preparation areas. Alcohol based hand products are not effective. Soap and water must be used
Fifth Disease	None	Good handwashing and hygiene; proper disinfection of changing tables, surfaces, and toys.
Hand-foot-and – mouth disease	None, unless fever present or child maintain hygiene or avoid close contact with others. Lesions must be scab over before return. Fever free without fever reducing medication for 24 hours	Good handwashing and hygiene; proper disinfection of changing tables, surfaces, and toys.

Impetigo	Exclude until after initiation of appropriate antibiotic treatment and lesion are covered or crusted.	Keep fingernails clean and short. Good handwashing and hygiene; proper disinfection of changing tables, surfaces, and toys.
Lice	Exclude until treatment or removal of live lice.	Avoid sharing or storing together personal items such as headgear, combs, and clothing.
Measles (Rubeola, measles virus)	Exclude for 4 days after rash onset; individual from day 7 through day 21 following earliest exposure. **At risk (see below)**	Assess exposure risk to who are at risk**; provide immunization record of exposed individuals to public health officials. Contacts without documented immunity (2 doses of vaccine or laboratory proof of immunity) should be vaccinated.
Ringworm	Exclude until treatment is initiated or lesions are covered.	Avoid direct contact, avoid sharing combs, brushes, hats, and clothing. Proper disinfection of surfaces and toys with a fungicidal agent.
Scabies	Exclude until treatment is complete.	Itching may continue for several weeks following treatment. It is not an indicator that treatment failed.
Varicella (chicken pox)	Exclude until lesions have dried and crusted; exclusion of at risk** usually not mandated, families should be notified of risk. Fever free without fever reducing medication for 24 hours	Assess exposure risk to who are at risk**; provide immunization record of exposed individuals to public health officials.
Meningitis (viral)	Excluded until fever has resolved for 24 hours. Fever free without fever reducing medication for 24 hours	Proper disinfection of surfaces with soap, water, and bleach-containing solution; treatment of contacts not necessary, no specific treatment.
Coronavirus (COVID-19)	Exclude until fever resolved for 24 hours Fever free without fever reducing medication for 24 hours	Handwashing and good personal hygiene including covering coughs and sneezes.

****At Risk = Person who are not immunized, with compromised immune system, or pregnant****

REVISED: May 14, 2015

APPROVED: June 15, 2015

APPROVED: August 13, 2020

REVISED: January 14, 2021

APPROVED: February 11, 2021

REVISED: September 9, 2021

APPROVED: October 14, 2021

REVISED: October 13, 2022

APPROVED: November 10, 2022

REVISED: December 14, 2023

APPROVED: January 11, 2024